

Use of passive air samplers in South America: a balance between global and local sources of Persistent Organic Pollutants (POPs)

Coordinator: Gilberto Fillmann (Institute of Oceanography, University Federal do Rio Grande, Brazil)

Researcher PUCP: Nadia Gamboa

Funded by: O Conselho Nacional de Desenvolvimento Científico e Tecnológico -CNPq, Brasil Programa Sul-Americano de Apoio às Atividades de Cooperação em Ciência e Tecnologia - PROSUL, **Rede Latino Americana de Monitoramento Atmosférico Passivo de POPs (LAPAN)**

Institutions involved: Institute of Oceanography(de dónde?), Universidade Federal do Rio Grande(do Sul?), Brazil/ Fundacão Oswaldo Cruz, Brazil/ Universidad (Nacional?) de Mar del Plata, Argentina / Centro EULA, Universidad de Concepcion, Chile/ Universidad de Cartagena, Colombia / Instituto de Ciencias Químicas y Ambientales -ESPOL, Ecuador/ Ministry of Environment of Ecuador/ Pontificia Universidad Católica del Perú/ Universidad de la República, Uruguay/ National Institute of Ecology, Mexico/ Fundación SIMBIOSIS para el Desarrollo, Bolivia/ Universidad Central de Venezuela/ University of Toronto Scarborough, Canada

Department of Science - Chemistry Section

Nadia Gamboa^a, Karina S. Miglioranza^b Ricardo Barra^c, Boris G. Johnson^d, Carola Resabala^e, Gabriela Eguren^f, Gary R. Anze^g, Raiza Fernandez^h, Frank Waniaⁱ, Gilberto Fillmann^j

^a Chemistry Section, Department of Science, PUCP

^b Universidad de Mar del Plata, Argentina

^c Universidad de Concepción, Chile

^d Universidad de Cartagena, Colombia

^e Instituto de Ciencias Químicas y Ambientales -ESPOL, Ecuador

^f Universidad de la República, Uruguay

^g Fundación SIMBIOSIS para el Desarrollo, Bolivia

^h Universidad Central de Venezuela

ⁱ, University of Toronto Scarborough, Canada

^j Instituto de Oceanografía, Universidad Federal do Rio Grande, Brasil

At present, Persistent Organic Pollutants (POPs) are widely distributed, remain in the environment, bioaccumulate through the food chain and show toxic effects. In order to reduce these emissions, about 100 passive air samplers (PAS) will be installed to cover

latitudinal gradients (from Ecuador to Antarctica) and longitudinal gradients (on the west to the east coast), in urban, rural and remote South America and Antarctica.

The project proposes the establishment of permanent regional air monitoring network of polycyclic aromatic hydrocarbons and other persistent organic pollutants such as chlorinated pesticides, polychlorinated biphenyl (PCBs) and polybrominated diphenyl ethers (PBDEs) by passive samplers and (agujas de coníferas) pine needles as bioindicators. The advantages offered include low cost of network installation and maintenance, in addition to the significant increase in spatial and temporal distribution. In this way, contributing to global studies that monitor the effectiveness of decisions established by the Stockholm Convention is expected.

In Peru, the PUCP already installed passive samplers at the *Estación Meteorológica Hipólito Unáñue* (Hipólito Unáñue Weather Station), *Estación Científica de la PUCP en Tambopata* (the PUCP Research Station in Tambopata) and the *Centro Piloto de Crianza de Sajino de la Universidad Nacional de la Amazonía Peruana* (Pilot Center of Sajino Breeding of the National University of the Peruvian Amazon.) Likewise, working on identifying representing sampling points transected in Lima and other regions. Finally, as an activity derived from the project, the intercalibration between research laboratories for the analysis of these pollutants are being conducted.